

# AstroChemical Newsletter #40

March 2019

You can access the full abstracts by clicking the paper titles. Submit your abstracts before the 25th of each month for inclusion in the following newsletter.

## Abstracts

### Protoplanetary discs: sensitivity of the chemical composition to various model parameters

**Wakelam, V.; Chapillon, E.; Dutrey, A.; Guilloteau, S.; Iqbal, W.; Coutens, A.; Majumdar, L.**

Protoplanetary discs are challenging objects for astrochemical models due to strong density and temperature gradients and due to the UV photons 2D propagation. In this paper, we have studied the importance of several model parameters on the predicted column densities of observed species. We considered: (1) two-phase (gas and homogeneous grains) or three-phase (gas, surface, and bulk of grains) models, (2) several initial compositions, (3) grain growth and dust settling, and (4) several cosmic-ray ionization rates. Our main result is that dust settling is the most crucial parameter. Including this effect renders the computed column densities sensitive to all the other model parameters, except cosmic-ray ionization rate. In fact, we found almost no effect of this parameter for radii larger than 10 au (the minimum radius studied here) except for N<sub>2</sub>H<sup>+</sup>. We also compared all our models with all the column densities observed in the protoplanetary disc around DM Tau and were not able to reproduce all the observations despite the studied parameters. N<sub>2</sub>H<sup>+</sup> seems to be the most sensitive species. Its observation in protoplanetary discs at large radius could indicate enough N<sub>2</sub> in the gas phase (inhibited by the three-phase model, but boosted by the settling) and a low electron abundance (favoured by low C and S elemental abundances).

MNRAS, Volume 484, Issue 2, p.1563-1573

DOI: [10.1093/mnras/stz068](https://doi.org/10.1093/mnras/stz068)

Full-text URL: <https://arxiv.org/abs/1901.10191>

### Molecular tracers of radiative feedback in Orion (OMC-1). Widespread CH<sup>+</sup> (J=1-0), CO (10-9), HCN (6-5), and HCO<sup>+</sup> (6-5) emission

**J. R. Goicoechea, M. G. Santa-Maria, E. Bron, D. Teyssier, N. Marcelino, J. Cernicharo, S. Cuadrado**

Young massive stars regulate the physical conditions, ionization, and fate of their natal molecular cloud. It is important to find tracers that help quantifying the stellar feedback processes that take place at different spatial scales. We present  $\sim 85$  arcmin<sup>2</sup> velocity-resolved maps of several submm molecular lines toward the closest high-mass star-forming region, OMC-1. The observed rotational lines include probes of warm and dense molecular gas that are difficult to detect from ground-based telescopes: CH<sup>+</sup> (1-0), CO (10-9), HCO<sup>+</sup> (6-5), and HCN (6-5). These lines trace a thin but very extended layer of molecular gas at high thermal pressure,  $P_{\text{th}} \sim 1\text{e}7\text{-}1\text{e}9$  K/cm<sup>3</sup>, associated with

the FUV-irradiated surface of OMC-1. The intense FUV field - emerging from massive stars in the Trapezium cluster - heats, compresses and photoevaporates the cloud edge. It also triggers the formation of reactive molecules such as CH<sup>+</sup>. The CH<sup>+</sup> (1-0) emission spatially correlates with the flux of FUV photons impinging the cloud:  $G_0$  from  $1e3$  to  $1e5$ . This correlation is supported by isobaric PDR models in the parameter space  $P_{th}/G_0 \sim [5e3-8e4]$  K/cm<sup>3</sup> where many PDRs seem to lie. The CH<sup>+</sup> (1-0) emission correlates with the extended emission from vibrationally excited H<sub>2</sub>, and with that of [CII]158 $\mu$ m and CO 10-9, all emerging from FUV-irradiated gas. These correlations link the presence of CH<sup>+</sup> to the availability of C<sup>+</sup> ions and of FUV-pumped H<sub>2</sub>( $v>0$ ) molecules. The parsec-scale CH<sup>+</sup> emission and narrow-line ( $\Delta v \sim 3$  km/s) mid-J CO emission arises from extended PDRs and not from fast shocks. PDR line tracers are the smoking gun of the stellar feedback from young massive stars. The PDR component in OMC-1 represents 5 to 10% of the total gas mass, however, it dominates the emitted line luminosity. These results provide insights into the source of submm CH<sup>+</sup> and mid-J CO emission from distant star-forming galaxies.

2019, A&A., 622, 91

DOI: [10.1051/0004-6361/201834409](https://doi.org/10.1051/0004-6361/201834409)

Full-text URL: <https://arxiv.org/abs/1812.00821>

## **An ALMA view of CS and SiS around oxygen-rich AGB stars**

**T. Danilovich, A. M. S. Richards, A. I. Karakas, M. Van de Sande, L. Decin and F. De Ceuster**

We aim to determine the distributions of molecular SiS and CS in the circumstellar envelopes of oxygen-rich asymptotic giant branch stars and how these distributions differ between stars that lose mass at different rates. In this study, we analyse ALMA observations of SiS and CS emission lines for three oxygen-rich galactic AGB stars: IK Tau, with a moderately high mass-loss rate of  $5e-6$  M $\odot$ /yr, and W Hya and R Dor with low mass-loss rates of  $\sim 1e-7$  M $\odot$ /yr. These molecules are usually more abundant in carbon stars but the high sensitivity of ALMA allows us to detect their faint emission in the low mass-loss rate AGB stars. The high spatial resolution of ALMA also allows us to precisely determine the spatial distribution of these molecules in the circumstellar envelopes. We run radiative transfer models to calculate the molecular abundances and abundance distributions for each star. We find a spread of peak SiS abundances with  $\sim 1e-8$  for R Dor,  $\sim 1e-7$  for W Hya, and  $\sim 3e-6$  for IK Tau relative to H<sub>2</sub>. We find lower peak CS abundances of  $\sim 7e-9$  for R Dor,  $\sim 7e-8$  for W Hya, and  $\sim 4e-7$  for IK Tau, with some stratifications in the abundance distributions. For IK Tau, we also calculate abundances for the detected isotopologues: C<sup>34</sup>S, <sup>29</sup>SiS, <sup>30</sup>SiS, Si<sup>33</sup>S, Si<sup>34</sup>S, <sup>29</sup>Si<sup>34</sup>S, and <sup>30</sup>Si<sup>34</sup>S. Overall, the isotopic ratios we derive for IK Tau suggest a lower metallicity than solar.

MNRAS 484, 494–509 (2019)

DOI: [10.1093/mnras/stz002](https://doi.org/10.1093/mnras/stz002)

Full-text URL: <https://arxiv.org/abs/1901.00070>

## **Methyl cyanide (CH<sub>3</sub>CN) and propyne (CH<sub>3</sub>CCH) in the low-mass protostar IRAS 16293-2422**

**Andron, Inès; Gratier, Pierre; Majumdar, Liton; Vidal, Thomas H. G.; Coutens, Audrey; Loison, Jean-Christophe; Wakelam, Valentine**

Methyl cyanide (CH<sub>3</sub>CN) and propyne (CH<sub>3</sub>CCH) are two molecules commonly used as gas thermometers for interstellar gas. They are detected in several astrophysical

environments and in particular towards protostars. Using data of the low-mass protostar IRAS 16293-2422 obtained with the IRAM 30-m single-dish telescope, we constrained the origin of these two molecules in the envelope of the source. The line shape comparison and the results of a radiative transfer analysis both indicate that the emission of CH<sub>3</sub>CN arises from a warmer and inner region of the envelope than the CH<sub>3</sub>CCH emission. We compare the observational results with the predictions of a gas-grain chemical model. Our model predicts a peak abundance of CH<sub>3</sub>CCH in the gas-phase in the outer part of the envelope, at around 2000 au from the central star, which is relatively close to the emission size derived from the observations. The predicted CH<sub>3</sub>CN abundance only rises at the radius where the grain mantle ices evaporate, with an abundance similar to the one derived from the observations.

MNRAS, Volume 481, Issue 4, p.5651-5659

DOI: [10.1093/mnras/sty2680](https://doi.org/10.1093/mnras/sty2680)

Full-text URL: <https://arxiv.org/abs/1812.05416>

## Survey Observations to Study Chemical Evolution from High-Mass Starless Cores to High-Mass Protostellar Objects II. HC<sub>3</sub>N and N<sub>2</sub>H<sup>+</sup>

**Kotomi Taniguchi, Masao Saito, T. K. Sridharan, Tetsuhiro Minamidani**

We have carried out survey observations of molecular emission lines from HC<sub>3</sub>N, N<sub>2</sub>H<sup>+</sup>, CCS, and cyclic-C<sub>3</sub>H<sub>2</sub> in the 81–94 GHz band toward 17 high-mass starless cores (HMSCs) and 28 high-mass protostellar objects (HMPOs) with the Nobeyama 45-m radio telescope. We have detected N<sub>2</sub>H<sup>+</sup> in all of the target sources except one and HC<sub>3</sub>N in 14 HMSCs and in 26 HMPOs. We investigate the N(N<sub>2</sub>H<sup>+</sup>)/N(HC<sub>3</sub>N) column density ratio as a chemical evolutionary indicator of massive cores. Using the Kolmogorov-Smirnov (K-S) test and Welch's t test, we confirm that the N(N<sub>2</sub>H<sup>+</sup>)/N(HC<sub>3</sub>N) ratio decreases from HMSCs to HMPOs. This tendency in high-mass star-forming regions is opposite to that in low-mass star-forming regions. Furthermore, we found that the detection rates of carbon-chain species (HC<sub>3</sub>N, HC<sub>5</sub>N, and CCS) in HMPOs are different from those in low-mass protostars. The detection rates of cyanopolyynes (HC<sub>3</sub>N and HC<sub>5</sub>N) are higher and that of CCS is lower in high-mass protostars, compared to low-mass protostars. We discuss a possible interpretation for these differences.

2019, The Astrophysical Journal, Volume 872, Number 2

DOI: [10.3847/1538-4357/ab001e](https://doi.org/10.3847/1538-4357/ab001e)

Full-text URL: <https://arxiv.org/abs/1901.06446>

## Electronic Transitions Responsible for C<sub>60</sub><sup>+</sup> Diffuse Interstellar Bands

**A. O. Lykhin, S. Ahmadvand, S. A. Varganov**

Despite the fact that C<sub>60</sub><sup>+</sup> recently became the first confirmed carrier of several diffuse interstellar bands (DIBs), the nature of the corresponding transitions is not understood. Using electronic structure methods, we show that the two strong C<sub>60</sub><sup>+</sup> DIBs cannot be explained by electronic transitions to the two different excited 2E<sub>1g</sub> states or the two spin-orbit components of the lowest 2E<sub>1g</sub> state, as suggested before. We argue that the strong DIBs at 9632 and 9577 Å correspond to the cold excitations from the non-Franck-Condon region of the ground electronic state to the two components of the lowest 2E<sub>1g</sub> state split by Jahn-Teller distortion. The weak DIBs at 9428 and 9365 Å are assigned to the first vibronic transitions involving the low-

energy vibrational modes and components of the lowest 2E1g electronic state.

J. Phys. Chem. Lett., 2019, 10 (1), pp 115-120

DOI: [10.1021/acs.jpcllett.8b03534](https://doi.org/10.1021/acs.jpcllett.8b03534)

Full-text URL: <https://pubs.acs.org/doi/full/10.1021/acs.jpcllett.8b03534>

## Organic molecules in the protoplanetary disk of DG Tau revealed by ALMA

**L. Podio, F. Bacciotti, D. Fedele, C. Favre, C. Codella, K. L. J. Rygl, I. Kamp, G. Guidi, E. Bianchi, C. Ceccarelli, D. Coffey, A. Garufi, and L. Testi**

Planets form in protoplanetary disks and inherit their chemical compositions. It is thus crucial to map the distribution and investigate the formation of simple organics, such as formaldehyde and methanol, in protoplanetary disks. We analyze ALMA observations of the nearby disk-jet system around the T Tauri star DG Tau in the o-H<sub>2</sub>CO 3(1,2)-2(1,1) and CH<sub>3</sub>OH 3(-2,2)-4(-1,4) E, 5(0,5)-4(0,4) A transitions at an unprecedented resolution of  $\sim 0.15''$ , i.e.,  $\sim 18$  au at a distance of 121 pc. The H<sub>2</sub>CO emission originates from a rotating ring extending from  $\sim 40$  au with a peak at  $\sim 62$  au, i.e., at the edge of the 1.3 mm dust continuum. CH<sub>3</sub>OH emission is not detected down to an r.m.s. of 3 mJy/beam in the 0.162 km/s channel. Assuming an ortho-to-para ratio of 1.8-2.8 the ring- and disk-height-averaged H<sub>2</sub>CO column density is  $\sim 0.3-4e14$  cm<sup>-2</sup>, while that of CH<sub>3</sub>OH is  $< 0.04-0.7e14$  cm<sup>-2</sup>. In the inner 40 au no o-H<sub>2</sub>CO emission is detected with an upper limit on its beam-averaged column density of  $\sim 0.5-6e13$  cm<sup>-2</sup>. The H<sub>2</sub>CO ring in the disk of DG Tau is located beyond the CO iceline (R(CO) $\sim 30$  au). This suggests that the H<sub>2</sub>CO abundance is enhanced in the outer disk due to formation on grain surfaces by the hydrogenation of CO ice. The emission peak at the edge of the mm dust continuum may be due to enhanced desorption of H<sub>2</sub>CO in the gas phase caused by increased UV penetration and/or temperature inversion. The CH<sub>3</sub>OH/H<sub>2</sub>CO abundance ratio is  $< 1$ , in agreement with disk chemistry models. The inner edge of the H<sub>2</sub>CO ring coincides with the radius where the polarization of the dust continuum changes orientation, hinting at a tight link between the H<sub>2</sub>CO chemistry and the dust properties in the outer disk and at the possible presence of substructures in the dust distribution.

8 pages, accepted on A&A Letters

DOI: [10.1051/0004-6361/201834475](https://doi.org/10.1051/0004-6361/201834475)

Full-text URL: <https://arxiv.org/abs/1902.02720>

## Statistical study of uncertainties in the diffusion rate of species on interstellar ice and its impact on chemical model predictions

**Iqbal, Wasim; Wakelam, Valentine; Gratier, Pierre**

Context. Diffusion of species on the dust surface is a key process for determining the chemical composition of interstellar ices. On the dust surface, adsorbed species diffuse from one potential well to another and react with other adsorbed reactants, resulting in the formation of simple and complex molecules. Aims: We study the impact on the abundances of the species simulated by the chemical codes by considering the uncertainties in the diffusion energy of adsorbed species. We aim to limit the uncertainties in the abundances as calculated by chemical codes by identifying the surface species that result in a larger error because of the uncertainties in their diffusion energy. Methods: We ran various cases with 2000-10 000 simulations in each case and varied the diffusion energies of some or all surface species randomly. We

calculated Pearson correlation coefficients between the abundances and the ratio of diffusion to binding energy of adsorbed species. We identified the species that introduce maximum uncertainty in the ice and gas-phase abundances. With these species we ran three sets, with 2000 simulations in each, to quantify the uncertainties they introduce. Results: We present the abundances of various molecules in the gas phase and also on the dust surface at different time intervals during the simulation. We show which species produce a large uncertainty in the abundances. We sorted species into different groups in accordance with their importance in propagating uncertainty in the chemical network. Conclusions: We show that CO, H<sub>2</sub>, O, N, and CH<sub>3</sub> are the key species for uncertainties in the abundances, while CH<sub>2</sub>, HCO, S and O<sub>2</sub> come next, followed by NO, HS, and CH. We also show that by limiting the uncertainties in the ratio of diffusion to binding energy of these species, we can eliminate the uncertainties in the gas-phase abundances of almost all the species.

A&A, Volume 620, December 2018

DOI: [10.1051/0004-6361/201833804](https://doi.org/10.1051/0004-6361/201833804)

Full-text URL: <https://arxiv.org/abs/1811.03488>

## ALMA Autocorrelation Spectroscopy of Comets: The HCN/H<sup>13</sup>CN Ratio in C/2012 S<sub>1</sub> (ISON)

**M. A. Cordiner, M. Y. Palmer, M. de Val-Borro, S. B. Charnley, L. Paganini, G. Villanueva, D. Bockelée-Morvan, N. Biver, A. Remijan, Y.-J Kuan, S. N. Milam, J. Crovisier, D. C. Lis, M. J. Mumma**

The Atacama Large Millimeter/submillimeter Array (ALMA) is a powerful tool for high-resolution mapping of comets, but the main interferometer (comprised of 50 × 12 m antennas) is insensitive to the largest coma scales due to a lack of very short baselines. In this Letter, we present a new technique employing ALMA autocorrelation data (obtained simultaneously with the interferometric observations), effectively treating the entire 12 m array as a collection of single-dish telescopes. Using combined autocorrelation spectra from 28 active antennas, we recovered extended HCN coma emission from comet C/2012 S<sub>1</sub> (ISON), resulting in a fourteen-fold increase in detected line brightness compared with the interferometer. This resulted in the first detection of rotational emission from H<sup>13</sup>CN in this comet. Using a detailed coma radiative transfer model accounting for optical depth and non-local thermodynamic equilibrium excitation effects, we obtained an H<sup>12</sup>CN/H<sup>13</sup>CN ratio of 88 ± 18, which matches the terrestrial value of 89. This is consistent with a lack of isotopic fractionation in HCN during comet formation in the protosolar accretion disk. The possibility of future discoveries in extended sources using autocorrelation spectroscopy from the main ALMA array is thus demonstrated.

2019, ApJ, 870, L26

DOI: [10.3847/2041-8213/aafb05](https://doi.org/10.3847/2041-8213/aafb05)

Full-text URL: <https://arxiv.org/abs/1901.08676>

## Announcements

### **Celebrating the First 40 Years of Alexander Tielens' Contribution to Science: the Physics and Chemistry of the ISM**

<https://tielens2019.sciencesconf.org/> Venue The symposium will be held 2—6

September 2019 in the historical Congres du Palais des Papes, Avignon, France

(<http://www.avignon-congres.com/>). Rational Xander Tielens has been driving research

in the fields of interstellar physics and chemistry and the cosmic cycle of matter with outstanding contributions for 40 years. With this meeting, we wish to celebrate his scientific achievements and discuss future research directions opened up by his contributions. The meeting will focus on the fields strongly influenced by Xander involving the physical and chemical processes that control the interstellar medium and its life cycle: PDRs, interstellar and circumstellar dust, PAHs, ices and astrochemistry. We will especially emphasize future opportunities offered by the powerful telescopes at our disposal such as, for example, ALMA, SOFIA, and JWST. The meeting will consist of invited reviews, invited and contributed talks, and posters. Key dates February 11: Opening of the registration and abstract submission on the symposium website June 1: Deadline of Abstract submission for oral contributions June 15: Announcement of the selected oral contributions June 20: Deadline of registration Please note that the symposium participation is restricted to 120 persons, based on first-signed first-selected, so do not delay your registration! Confirmed invited speakers L. Allamandola, N. Balucani, A. Boogert, F. Boulanger, S. Cazaux, J. Cernicharo, L. d'Hendecourt, T. de Graauw, J. Goicoechea, H. Habing, D. Hollenbach, C. Joblin, B. Lefloch, M. Kaufman, C. Kemper, M. Meixner, T. Millar, T. Onaka, M-E. Palumbo, E. Roueff, K. Schuster, E. van Dishoeck, R. Waters Scientific Organizing Committee C. Ceccarelli (chair), A. Candian (co-chair), J. Cami, C. Dominik, L. Hornekaer, K. Justtanont, E. Peeters, M. Wolfire Local Organizing Committee B. Lefloch (chair), E. Bianchi, C. Borye, M. Bouvier, M. De Simone, M-H Sztefek Contacts loc-XT2019@univ-grenoble-alpes.fr

## **Symposium “The Periodic Table Through Space and Time” September 9-13 of 2019, Saint-Petersburg (Russia)**

The 2019 year is claimed as the International Year of the Periodic Table of Chemical Elements. This initiative was supported by many international organizations, including the International Astronomical Union (IAU). In September 9-13 of 2019, Saint-Petersburg (Russia) hosts the XXI Jubilee Mendeleev Congress on General and Applied Chemistry (<http://mendeleev2019.ru/index.php/en/>). During this Congress, in Sept. 10-13 of 2019, we hold the Symposium “The Periodic Table Through Space and Time”. Main topics of the Symposium are: - Big Bang and stellar nucleosynthesis - Chemical evolution of the Universe: observations and models - Astrochemistry - Elemental abundances: a key to stellar physics Invited speakers: to be confirmed. Scientific Organizing Committee: Ewine van DISHOECK (The Netherlands, Co-Chair), Boris SHUSTOV (Co-Chair, Russia), Lyudmila MASHONKINA (Russia, Scientific Secretary), Sergey BLINNIKOV (Russia), John COWAN (USA), Charles Cowley (USA), Alexander DOLGOV (Russia), Nicolas GREVESSE (Belgium), John LANDSTREET (Canada), Alexander LUTOVINOV (Russia), Ken'ichi NOMOTO (Japan), Nikos PRANTZOS (France), Friedrich-Karl THIELEMANN (Switzerland), Dmitry WIEBE (Russia), Gang ZHAO (China). On behalf of the Organizing Committee of Congress and the SOC of the Symposium we are pleased to invite academic and university researchers, postgraduate students, and students to participate in the Symposium “The Periodic Table Through Space and Time”. The online registration and abstracts submission is now opened! The guidelines and abstract template can be found on the Congress website page <http://mendeleev2019.ru/index.php/en/about-the-event-3/abstracts> Please note that timely registration is essential to ensure that visa formalities can be completed in due time. For questions related to the Symposium, you may contact us via e-mail: [mendeleev150@inasan.ru](mailto:mendeleev150@inasan.ru)

## **Postdoc position available for Doctors in Chemistry or Physics**

A Postdoc Position is available at University of Castilla-La Mancha (UCLM). Department of Physical Chemistry. Ciudad Real (Spain). The postdoc is funded by the European project: Gas and Dust from the Stars to the Laboratory: Exploring the NANOCOSMOS

(ERC-2013-SyG Grant Agreement nº 610256). <http://www.icmm.csic.es/nanocosmos/>  
The main objective is: Measurement of Rate Coefficients of Radical-Molecule Reactions in the Gas-phase At Ultra-low Temperatures Using A Pulsed CRESU (Uniform Supersonic Flow) System Coupled to Pulsed Laser Photolysis and Laser Induced Fluorescence. Duration is up to July 2020. Preferred experience: spectroscopy, gases, lasers, programming. Send your CV and two reference letters to Dr. E. Jiménez (Elena.Jimenez@uclm.es).